

Head Injury Report



(Name of child) _____ has sustained a head injury at Milton Mount Playgroup on _____ at approximately _____ and has been monitored since the accident and we have not identified anything that caused concern up to the time of them going home.

Details _____

If any of these symptoms are present, particularly loss of consciousness (even for a short period of time), you should call an emergency ambulance (999 / 112) or NHS Direct on 111 -Lasting headache that gets worse or is still present over six hours after the injury;

- Extreme difficulty in staying awake, or still being sleepy several hours after the injury. It is fine to let children go to sleep after a slight bump to the head, but you should check on them regularly and make sure you are able to wake them.
- Nausea and vomiting several hours after the injury;
- Unconsciousness or coma;
- Unequal pupil size;
- Confusion, feeling lost or dizzy, or difficulty making sense when talking;
- Pale yellow fluid or watery blood, coming from the ears or nose (this suggests a skull fracture);
- Bleeding from the scalp that cannot be quickly stopped;
- Not being able to use part of the body, such as weakness in an arm or leg;
- Difficulty seeing or double vision;
- Slurred speech; and
- Having a seizure or fit.

Head Injury Report



(Name of child) _____ has sustained a head injury at Milton Mount Playgroup on _____ at approximately _____ and has been monitored since the accident and we have not identified anything that caused concern up to the time of them going home.

Details _____

If any of these symptoms are present, particularly loss of consciousness (even for a short period of time), you should call an emergency ambulance (999 / 112) or NHS Direct on 111 -Lasting headache that gets worse or is still present over six hours after the injury;

- Extreme difficulty in staying awake, or still being sleepy several hours after the injury. It is fine to let children go to sleep after a slight bump to the head, but you should check on them regularly and make sure you are able to wake them.
- Nausea and vomiting several hours after the injury;
- Unconsciousness or coma;
- Unequal pupil size;
- Confusion, feeling lost or dizzy, or difficulty making sense when talking;
- Pale yellow fluid or watery blood, coming from the ears or nose (this suggests a skull fracture);
- Bleeding from the scalp that cannot be quickly stopped;
- Not being able to use part of the body, such as weakness in an arm or leg;
- Difficulty seeing or double vision;
- Slurred speech; and
- Having a seizure or fit.